

United Healthcare VBP Measures – 2021

Medicaid Line of Business

Code	Measure	Description
SAA	Adherence to Antipsychotic Medications for Individuals With Schizophrenia	Percentage of members ages 18 and older during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80 percent of the treatment period.
BCS	Breast Cancer Screening	Percentage of female members ages 50–74 who had a mammogram screening Oct. 1 two years prior to the measurement year through Dec. 31 of the measurement year.
CCS	Cervical Cancer Screening	Percentage of female members ages 21–64 who were screened for cervical cancer using either of the following criteria: <ul style="list-style-type: none"> •Women ages 21–64 who had cervical cytology performed in the measurement year or two years prior •Women ages 30–64 who had cervical cytology/high-risk human papillomavirus (hrHPV) performed in the measurement year or 4 years prior. The woman must have been at least 30 on the date of the test •Women ages 30–64 who had cervical high-risk human papillomavirus (hrHPV) testing performed in the measurement year or 4 years prior
COL	Colorectal Cancer Screening	Percentage of members ages 50–75 who had one or more appropriate screenings for colorectal cancer. <p>FOBT: Current calendar year</p> <p>FIT-DNA: Current calendar year or two years prior</p> <p>Flex Sig: Current calendar year to previous 4 calendar years</p> <p>Colonoscopy: Current calendar year to previous 9 calendar years</p>
CDC	Comprehensive Diabetes Care: Eye Exam (Retinal) Performed	Percentage of members ages 18–75 with diabetes (Types 1 and 2) who had any one of the following: <ul style="list-style-type: none"> •Retinal or dilated eye exam by an optometrist or ophthalmologist in the measurement year •Negative retinal or dilated eye exam by an optometrist or ophthalmologist in the year prior to the measurement year •Bilateral eye enucleations any time during their history through Dec. 31 of the measurement year
IMA	Immunizations for Adolescents: Combination 2	Percentage of adolescents age 13 who had one dose of meningococcal vaccine, one tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine, and completed the human papillomavirus (HPV) vaccine series by their 13th birthday.
PPC	Postpartum Care	Deliveries of live births on or between Oct. 8 of the year prior to the measurement year and Oct. 7 of the measurement year and percentage of women who had a live birth that had a postpartum visit on or between 7–84 days after delivery.
WCV	Child and Adolescent Well Care Visits	Percentage of members ages 3–21 who had one or more well-child visits with a primary care provider or OB-GYN during the measurement year.
W30	Well Child Visits in the First 30 Months of Life	Percentage of members who turned 15–30 months old during the measurement year and had the recommended number of well-child visits with a primary care provider. <ul style="list-style-type: none"> •Children who turned 15 months old during the measurement year: Six or more well-child visits. •Children who turned 30 months old during the measurement year: Two or more well-child visits.
FUA	Follow-Up After Emergency	The percentage of ED visits for members ages 13 and older with a principal

	Department Visit for Alcohol and Other Drug Dependence (7 Days)*	<p>diagnosis of alcohol or other drug abuse or dependence (AOD) and who had a follow-up visit for AOD.</p> <ul style="list-style-type: none"> •The percentage of ED visits for which the member received follow-up for AOD within the 7 days after the visit (8 days total)
VLS	HIV Viral Load Suppression*	The percentage of patients, regardless of age, with a diagnosis of HIV with a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.